

- 28 Gan WQ, Mannino DM, Jemal A. Socioeconomic disparities in secondhand smoke exposure among US never-smoking adults: the National Health and Nutrition Examination Survey 1988–2010. *Tob Control* 2015;24:568–73.
- 29 Aurrekoetxea JJ, Murcia M, Rebagliato M, et al. Factors associated with second-hand smoke exposure in non-smoking pregnant women in Spain: self-reported exposure and urinary cotinine levels. *Sci Total Environ* 2014;470–1:1189–96.
- 30 Kuntz B, Lampert T. Social disparities in parental smoking and young children's exposure to secondhand smoke at home: a time-trend analysis of repeated cross-sectional data from the German KiGGS study between 2003–2006 and 2009–2012. *BMC Public Health* 2016;16:485.
- 31 Protano C, Valeriani F, Macedonio A, et al. Family-based social determinants and child health: cross-sectional study. *Pediatr Int* 2017;59:201–8.
- 32 Arechavala T, Continente X, Pérez-Ríos M, et al. Validity of self-reported indicators to assess secondhand smoke exposure in the home. *Environ Res* 2018;164:340–5.

.....  
*The European Journal of Public Health*, Vol. 29, No. 5, 849–855

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Public Health Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

doi:10.1093/eurpub/ckz076 Advance Access published on 23 April 2019

## Inequity in postpartum healthcare provision at home and its association with subsequent healthcare expenditure

Jacqueline Lagendijk <sup>1</sup>, Eric A.P. Steegers<sup>1</sup>, Jasper V. Been<sup>1,2,3</sup>

1 Department of Obstetrics and Gynaecology, Erasmus MC, University Medical Centre Rotterdam, Rotterdam, the Netherlands

2 Division of Neonatology, Department of Paediatrics, Erasmus MC – Sophia Children's Hospital, University Medical Centre Rotterdam, Rotterdam, the Netherlands

3 Department of Public Health, Erasmus MC, University Medical Centre Rotterdam, Rotterdam, the Netherlands

**Correspondence:** Jacqueline Lagendijk, Department of Obstetrics and Gynaecology, Erasmus MC, University Medical Centre Rotterdam, PO Box 2040, 3000 CA Rotterdam, The Netherlands, Tel: +31 (0) 10 703 68 86, Fax: +31 (0) 10 703 68 15, e-mail: j.lagendijk.2@erasmusmc.nl

**Background:** Provision of postpartum care can support new families in adapting to a new situation. We aimed to determine whether various determinants of socioeconomic status (SES) were associated with utilization of postpartum care. In addition, to stress the relevance of increasing postpartum care uptake among low SES-groups, an assessment of the potential (cost-)effectiveness of postpartum care is required. **Methods:** National retrospective cohort study using linked routinely collected healthcare data from all registered singleton deliveries (2010–13) in the Netherlands. Small-for-gestational age and preterm babies were excluded. The associations between SES and postpartum care uptake, and between uptake and health care expenditure were studied using multivariable regression analyses. **Results:** Of all 569 921 deliveries included, 1.2% did not receive postpartum care. Among women who did receive care, care duration was below the recommended minimum of 24 h in 15.3%. All indicators of low SES were independently associated with a lack in care uptake. Extremes of maternal age, single parenthood and being of non-Dutch origin were associated with reduced uptake independent of SES determinants. No uptake of postpartum care was associated with maternal healthcare expenses in the highest quartile: aOR 1.34 (95% CI 1.10–1.67). Uptake below the recommended amount was associated with higher maternal and infant healthcare expenses: aOR 1.09 (95% CI 1.03–1.18) and aOR 1.20 (95% CI 1.13–1.27), respectively. **Conclusion:** Although uptake was generally high, low SES women less often received postpartum care, this being associated with higher subsequent healthcare expenses. Strategies to effectively reduce these substantial inequities in early life are urgently needed.

### Introduction

The postpartum period is a critical transitional period not only for babies but also in the lives of new mothers.<sup>1</sup> Adequate care provision during this period by skilled maternity care professionals enables an optimal start for the new family. A healthy start following childbirth may be of substantial short and long term benefit for maternal and child wellbeing, and as such has the potential to reduce healthcare associated costs.<sup>2,3</sup>

The uptake of healthcare overall and the incidence of adverse health outcomes during the postpartum period are closely linked to different determinants of one's socioeconomic position; persons with a lower socioeconomic position tend to make less use of

routine or preventive healthcare,<sup>4,5</sup> and have a higher incidence of adverse health outcomes.<sup>3,6–10</sup> Although a number of studies examined this relationship, the association between SES and use of postpartum care has not been investigated previously.

The strong position of primary care in the Netherlands, which includes easy access to postpartum care at home during the early postpartum period (figure 1), provides considerable potential to promote equity in maternal and infant health. This study seeks to describe the patterns of utilization of postpartum care services using a national population-based study, assessing: (i) whether different determinants of SES—represented by individual level, household level and area-level indicators—were associated with uptake of postpartum care and (ii) whether any inequalities translated in

**Box 1. Postpartum care in the Netherlands**

In the Netherlands, postpartum care professionals provide care in a family's home during the postpartum period. This care is initially delivered for six to eight hours a day, tapering to fewer hours per day towards the end of the care period, which usually consist of eight consecutive days following childbirth. The purpose of this postpartum care is to provide maternal and infant care by giving informational and emotional support, by providing comprehensive care for the mother and child dyad, and by enhancing maternal empowerment.

The duration of postpartum care is indexed in advance for each woman individually during a scheduled home visit during the second or third trimester of pregnancy.

The minimal amount of postpartum care is set at 24 hours (i.e. three hours per day for eight consecutive days after childbirth). The recommended amount of care is set at 49 hours, and the maximum amount of care is 80 hours depending on specific indications, for example an (imminent) mastitis or symptoms of psychological illness. When a mother and child dyad is hospitalised after childbirth (for example when an infant is born preterm or SGA) the indexed amount of postpartum care at home is reduced for every day that they remain in the hospital.

Skilled nurses who have a lower secondary education degree for postpartum care provide this care.

The costs for postpartum care are covered within the basic package of health insurance, which is obligatory for Dutch citizens, with exception of an out-of-pocket payment per hour of care received (i.e. € 4.30 in 2018) (<http://www.digitalezorg.nl/digitale/uploads/2015/03/netherlands.pdf>).

**Figure 1** Postpartum care in the Netherlands

uptake of care translated into differences in subsequent healthcare expenditures for mother and child in the first year after childbirth, as an estimate of potential (cost-)effectiveness.

## Methods

We conducted a national population-based retrospective cohort study of women living in the Netherlands who delivered a live singleton baby between 1 January 2010 and 31 December 2013. Routinely collected healthcare and claims data were linked at the individual level across various national databases. First, we studied the association between different determinants of low SES and the uptake of postpartum care. Second, we studied the association between the uptake of postpartum care and healthcare expenditures for mother and child in the following year. We used the RECORD statement to guide reporting of our findings (Supplementary file S1).<sup>11</sup>

### Study design and setting

Population-based retrospective cohort study from 1 January 2010 through 31 December 2013 using routinely collected healthcare data from Statistics Netherlands (translated Dutch name: 'Central Bureau of Statistics', abbreviation 'CBS').

### Participants

All registered pregnancies among women living in the Netherlands who delivered a live singleton baby at 24 or more completed gestational weeks between 2010 and 2013.

### Exposure variables

For the first part of this study, multiple determinants of SES including individual, household and area-level SES indicators constituted the exposures of interest. Disposable household income was used as an individual SES indicator, defined as the sum available from the household income for final consumption and savings (i.e. net income) and divided into quartiles. Mother's highest educational qualification, based on the International Standard Classification of Education (ISCED) (<http://uis.unesco.org/sites/default/files/documents/international-standard-classification-of-education-isced-2011-en.pdf>), was considered a second individual SES indicator. Three groups were considered: lower

education (pre-primary, primary and lower secondary education), intermediate education (upper secondary education, post-secondary non-tertiary education) and higher education (first stage of tertiary education, second stage of tertiary education).

Home ownership was considered a household indicator of SES and was dichotomized into owner-occupiers and no-owners (i.e. renters and others).

Neighbourhood deprivation was considered an area-level SES indicator and was based on the Neighbourhood Deprivation Index (NDI) formulated by NIVEL in 2012.<sup>12</sup> Deprivation was defined at an NDI of 5.5% (i.e. 885 000 people).

In the second part, the exposure was the uptake of any postpartum care, and—in a secondary analysis—the uptake of postpartum care above the recommended minimum (i.e. 24 h) among those who did receive postpartum care.

### Covariates

Covariates were selected based on their association with the outcome variables or both the outcome and the exposure variables: maternal age, parity, country of origin, parenthood household status, level of urbanization and small-for-gestational age and preterm babies. Details are presented in Supplementary file S2.

### Outcomes

#### Determinants of low SES and uptake of postpartum care

Uptake of postpartum care was derived from data regarding healthcare expenditures. Expenditures were provided per annum; therefore pregnancies from women who gave birth more than once within 1 year were excluded for all analysis. The amount of postpartum care was calculated by dividing the total postpartum care expenditures within 1 year by the eligible compensation per hour of care, which differed per year.<sup>13</sup> Uptake of postpartum care was dichotomized into 'No', and 'Yes' (any amount of postpartum care). The secondary outcome was postpartum care uptake above the minimum (i.e. 24 h), as assessed among all women who did make use of postpartum care. The uptake of the minimum amount of care was dichotomized.

## Uptake of postpartum care and healthcare expenditures

Annual total healthcare expenditures were obtained separately for mother and child. Quartiles of annual healthcare expenditures were formed for each and dichotomized into 'low' (expenses within the first three quartiles) and 'high' (expenses in the fourth quartile). Healthcare expenditure data were only available at an aggregated level per annum. We were therefore able to reliably evaluate health care costs in the year post-delivery only among those women delivering close to closing of the year. As such, we pragmatically considered total healthcare costs in the subsequent year following delivery in December a reasonable estimate of healthcare expenditure in the year post-delivery, and excluded deliveries in January to November from these analyses. Healthcare expenditures are subdivided based on a combination of diagnosis and treatment codes enabling us to exclude all healthcare expenses that were labelled as pregnancy-related. In addition, we excluded women with more than one pregnancy during the study period (i.e. 2010–13) because having consecutive pregnancies over a 2-year period could influence healthcare expenditures at the annual level.

### Data sources and linkage

The available data for this study were linked across different national registries by CBS using the unique citizen service number (BSN) or the identification number of the Dutch Population Register (Dutch: A-number). Linkage with this information is feasible in 98–100% of all procedures undertaken by CBS. Details about the individual-level linkage across various routinely collected datasets are presented in Supplementary file S2.

### Potential for bias

The data in this study are based on routinely collected healthcare data. There was a reasonably high proportion of missing values in some registries that could have introduced different biases. We applied multiple imputation using chained equations to account for this missing data in baseline characteristics. Multiple predictor variables were included to inform the multiple imputation process, forming 10 datasets. Results across the sets were combined using Rubin's Rules.<sup>14</sup>

### Statistical methods

We analysed the two associations under study using logistic regression analysis.

Infants born preterm or SGA and their mothers tend to remain in the hospital during most of the time that the mothers would otherwise be amenable to receiving postpartum care in the home situation (figure 1). Therefore, we excluded deliveries with these outcomes for all analyses because postpartum care uptake would otherwise be underestimated due to prolonged hospital admission.

### Determinants of low SES and the uptake of postpartum care

The association between various determinants of low SES and postpartum care uptake (first), and uptake above the minimum (second) was analysed. All indicators of SES as exposure variables, and the predefined covariates were included in the analysis to minimize potential confounding.

### Uptake of postpartum care and healthcare expenditures

The second model analysed the association between postpartum care uptake and healthcare expenditures for mother and child. We

accounted for all SES indicators and all covariates included in the first model.

### Sensitivity analyses

Consecutive pregnancies within the same mother have more characteristics in common than pregnancies between women. To assess whether this dependency of data affected our findings, we reran the model that analyses the first association with additional accounting for clustering at the individual level.

To assess whether the multiple imputed data were biased, we reran the two models on complete cases only.

### Accessibility of protocol and programming code

Upon request all programming codes and the study protocol are available with the principal investigator.

### Details of ethics approval

According to Dutch law, formal ethical assessment of the study protocol was not needed as the study did not involve an intervention and data from CBS are anonymized [based on guidance from the Central Committee on Research Involving Human Subjects (WMO) and the Dutch Personal Data Protection Act].

CBS collects and produces population statistics, referred to as non-public microdata, for all registered Dutch citizens. Under strict conditions, these data are accessible for scientific research. The research board of CBS has reviewed and approved the study protocol (project number 7883).

## Results

### Participants

During the study period, 683 163 deliveries were registered with CBS. After applying the pre-specified exclusion criteria, the final sample included 569 921 deliveries (Supplementary figure S1). For investigation of the association between postpartum care uptake and healthcare expenditures, we additionally excluded deliveries in January through November, and consecutive pregnancies within individual women during the study period. The final sample for this analysis contained 44 458 deliveries (Supplementary figure S1).

### Determinants of low SES and uptake of postpartum care

#### Univariable associations

Table 1 presents the descriptive statistics for the study sample, by uptake of postpartum care. Of all deliveries included, 1.2% did not receive any postpartum care. Data on the uptake of postpartum care were missing for 4.8% of all deliveries. Women who did not use postpartum care were more often: multiparous (67.9% vs. 54.2%), single parents (20.1% vs. 7.7%), born outside the Netherlands (2.9% vs. 0.6%) and they more often lived in deprived neighbourhoods (19.1% vs. 6.8%; table 1). Among women who did receive postpartum care, care duration was below the recommended minimum of 24 h in 15.3% (Supplementary table S1). These deliveries were also associated with indicators of low SES when compared with deliveries with postpartum care uptake above the minimum amount (Supplementary table S1).

#### Multivariable associations

All indicators of low SES were consistently and strongly associated with no uptake of postpartum care after mutual adjustment (table 2). Similarly, among mothers who did receive postpartum care, low SES indicators were associated with care uptake below the minimum (table 2). Extremes of maternal age, single parenthood

**Table 1** Descriptive statistics of all deliveries by uptake of postpartum care (yes, no, missing)

Total population	Postpartum care uptake							
	<i>n</i> =569 921		<i>n</i> =535 470		<i>n</i> =6833		<i>n</i> =27 618	
		%		%		%		%
Maternal age								
<20	6837	1.2	6231	1.2	237	3.5	369	1.3
20–40	552 753	97.0	519 882	97.1	6325	92.6	26 546	96.1
>40	10 331	1.8	9357	1.7	271	4.0	703	2.5
Parity								
Primiparous	259 330	45.5	245 298	45.8	2196	32.1	11 836	42.9
Multiparous	310 591	54.5	290 172	54.2	4637	67.9	15 782	57.1
Country of origin								
The Netherlands	414 243	72.7	393 408	73.5	2544	37.2	18 291	66.2
Morocco	24 726	4.3	22 920	4.3	980	14.3	826	3.0
Turkey	18 985	3.3	17 989	3.4	515	7.5	481	1.7
Suriname	13 802	2.4	12 864	2.4	372	5.4	566	2.0
Netherlands Antilles	6503	1.1	5864	1.1	189	2.8	450	1.6
Other Non-Western	36 253	6.4	32 077	6.0	1216	17.8	2960	10.7
Other Western	55 409	9.7	50 348	9.4	1017	14.9	4044	14.6
Parenthood status								
Single parent	44 576	7.8	41 130	7.7	1377	20.2	2069	7.5
Two parents	521 140	91.4	491 138	91.7	5184	75.9	24 818	89.9
Other	4197	0.7	3195	0.6	272	4.0	730	2.6
Missing	8	0.0	7	0.0	0	0.0	1	0.0
Urbanized area								
Yes	163 610	28.7	155 696	29.1	1247	18.2	6667	24.1
No	406 311	71.3	379 774	70.9	5586	81.8	20 951	75.9
SES indicators								
Education								
Lower education	74 984	13.2	70 317	13.1	2052	30.0	2615	9.5
Intermediate education	163 305	28.7	156 371	29.2	1438	21.0	5496	19.9
Higher education	197 725	34.7	184 817	34.5	955	14.0	11 953	43.3
Missing	133 907	23.5	123 965	23.2	2388	34.9	7554	27.4
Low-disposable income								
Yes	131 290	23.0	122 207	22.8	3445	50.4	5638	20.4
No	416 580	73.1	393 035	73.4	2993	43.8	20 552	74.4
Missing	22 051	3.9	20 228	3.8	395	5.8	1428	5.2
Home ownership								
No-owners	146 307	25.7	135 846	25.4	3984	58.3	6477	23.5
Owner-occupiers	401 563	70.5	379 396	70.9	2454	35.9	19 713	71.4
Missing	22 051	3.9	20 228	3.8	395	5.8	1428	5.2
Neighbourhood deprivation								
Yes	39 526	6.9	36 248	6.8	1305	19.1	1973	7.1
No	530 395	93.1	499 222	93.2	5528	80.9	25 645	92.9

Values are presented as numbers and percentage.

and being of non-Dutch origin were associated with reduced uptake of postpartum care independent of individual and area-level SES.

### Sensitivity analyses

Consistent results were obtained in all sensitivity analyses for robustness checks, including those accounting for clustering of pregnancies at the individual level, and those analysing complete cases only (Supplementary table S2).

### Uptake of postpartum care and healthcare expenditures

#### Univariable associations

Descriptive statistics for the subgroup of 44 458 deliveries in December, were similar to those of all deliveries (Supplementary table S3). The prevalence of low SES indicators increased steadily across the four quartiles of maternal healthcare expenditure, with the highest quartile having the highest prevalence of low SES indicators: lowest educational level 23.0% in the highest quartile vs. 17.3% across the other quartiles, low disposable income 30.0% vs. 25.2%, no home-ownership 34.4% vs. 27.7% and living in a deprived neighbourhood 9.0% vs. 7.1% (Supplementary table S3). This tendency

was not seen across the four quartiles of infant healthcare expenditure, were the prevalence of low maternal SES indicators in the highest quartile was comparable with the prevalence across the other quartiles (data not presented). The percentage of women who did not receive postpartum care was highest in the fourth quartile of maternal healthcare expenses (2.1% in the highest quartile vs. 1.2% across the other quartiles; Supplementary table S3).

#### Multivariable associations

Not receiving postpartum care, or having postpartum care uptake below the minimum, was associated with a significantly higher odds of having maternal healthcare expenditure within the highest quartile in the year following child birth: aOR 1.34; 95% CI 1.10–1.67;  $P$  0.004, and aOR 1.09; 95% CI 1.03–1.18;  $P$  0.005, respectively (table 3). Deliveries followed by postpartum care uptake below the minimum were in addition associated with infant healthcare expenditure within the highest quartile in the first year after birth (aOR 1.20; 95% CI 1.13–1.27;  $P$  < 0.001) (table 3).

#### Sensitivity analyses

The association between no uptake of postpartum care and maternal healthcare expenses during the first year after childbirth was

**Table 2** Multivariable models of the association between SES indicators and 1) postpartum care uptake and 2) uptake above the minimum (i.e. 24 h)

	Postpartum care uptake (n = 569 921)		Uptake above the minimum amount (n = 535 470)	
	aOR (95% CI)	P value	aOR (95% CI)	P value
<b>Covariates</b>				
Maternal age				
<20 years	0.70 (0.61–0.81)	<0.001	0.57 (0.54–0.60)	<0.001
20–40 years (ref)	1		1	
>40 years	0.53 (0.47–0.60)	<0.001	0.78 (0.73–0.82)	<0.001
Parity				
Primiparous (ref)	1		1	
Multiparous	0.59 (0.55–0.62)	<0.001	0.99 (0.98–1.01)	0.601
Parenthood				
Single parent	0.82 (0.76–0.88)	<0.001	0.82 (0.80–0.84)	<0.001
Two parents (ref)	1		1	
Other	0.22 (0.19–0.25)	<0.001	0.40 (0.37–0.43)	<0.001
Country of origin				
The Netherlands (ref)	1		1	
Morocco	0.37 (0.34–0.40)	<0.001	0.23 (0.22–0.24)	<0.001
Turkey	0.44 (0.40–0.49)	<0.001	0.26 (0.25–0.27)	<0.001
Suriname	0.46 (0.41–0.51)	<0.001	0.29 (0.28–0.30)	<0.001
Net. Antilles	0.50 (0.43–0.58)	<0.001	0.32 (0.30–0.34)	<0.001
Other Non-Western	0.40 (0.37–0.43)	<0.001	0.22 (0.22–0.23)	<0.001
Other Western	0.41 (0.41–0.48)	<0.001	0.41 (0.40–0.42)	<0.001
<b>SES indicators</b>				
<b>Individual</b>				
Education				
Lower education	0.62 (0.57–0.66)	<0.001	0.65 (0.64–0.67)	<0.001
Inter. education (ref)	1		1	
Higher education	1.21 (1.12–1.32)	<0.001	1.39 (1.36–1.42)	<0.001
Low-disposable income				
Yes	0.72 (0.68–0.77)	<0.001	0.69 (0.68–0.71)	<0.001
<b>Household</b>				
Home ownership				
No-owners	0.55 (0.52–0.59)	<0.001	0.51 (0.50–0.52)	<0.001
Owner-occupiers (ref)	1		1	
<b>Area-level</b>				
Neighbourhood deprivation				
Yes	0.80 (0.75–0.85)	<0.001	0.79 (0.77–0.81)	<0.001

Presented are adjusted odds ratios (aOR) and their 95% confidence intervals (95%CI). All p-values are two-sided. Results for the uptake of care and the minimum uptake of care are presented separately.

consistent in the sensitivity analysis using complete cases only [aOR 1.54 (95% CI 1.23–1.92;  $P < 0.001$ )].

## Discussion

Using a national linked dataset of over half a million singleton pregnancies, we found that all indicators of low SES were associated with no uptake of postpartum care and with uptake of care below the recommended minimum. This lack of postpartum care uptake was associated with higher healthcare expenses in the first year after childbirth. For the first time, we demonstrate that postpartum care may be a cost-effective tool but is least provided to those who are most likely to benefit from it.

Strengths of this study include the very large and nationally representative sample and the use of a unique individual-level linkage across various routinely collected datasets of relevant medical and socioeconomic data. The relationship between low SES and lack of uptake of postpartum care was highly consistent across the various SES indicators. Although at the population level the proportion of women not receiving postpartum care is very small, we have shown that these women represent a marginalized group and may therefore benefit from efforts to improve their care.

In addition, associations between low SES and postpartum care uptake as well as between postpartum care uptake and subsequent healthcare expenditure showed a dose-response association. The largest differences were present between women who did not

receive postpartum care and those who received care above the minimum amount. The findings of both analyses were furthermore highly robust in sensitivity analyses. In the absence of major changes to the system used for indicating the amount of postpartum care and of the health care insurance system in the Netherlands, the data used in this study (from 2010 to 2013) may be considered generalizable to the current day.

Our study also has limitations. First, some of the national registries from Statistics Netherlands have a reasonably high percentage of missing values. For example, the percentage of missing values on a woman's highest educational qualification was as high as 30%. Upward educational-attainment biases could have influenced the registered data. To minimize bias within the imputed data, we had all SES indicators and outcome variables inform the imputation process. Sensitivity analyses on complete cases only showed similar results to the main analyses, supporting validity of the imputation and robustness of the findings. Second, we lacked information on medical conditions of women and infants. Having a medical condition that requires inpatient treatment could directly affect the uptake of postpartum care, as this care is provided only in the primary care setting (i.e. at home or in a primary care birth center). A third limitation is that the provided postpartum care is expressed in total expense rather than days of care received, making derivation necessary. In addition, we did not have information on the number of days spent in the hospital prior to receiving postpartum care, which may have biased our findings. Somewhat

**Table 3** Multivariable model of the association between the uptake of postpartum care and the lack in uptake and maternal and infant healthcare expenses within the highest quartile

	Maternal health care expenditures highest quartile		Infant health care expenditures highest quartile	
	aOR (95% CI)	P value	aOR (95% CI)	P value
Primary analyses				
Maternity care uptake (n=44 458)				
No	1.34 (1.10–1.67)	0.004	1.12 (0.94–1.35)	0.205
Yes (ref)	1		1	
Maternity care >24 h (n=41 583)				
No	1.09 (1.03–1.18)	0.005	1.20 (1.13–1.27)	<0.001
Yes (ref)	1		1	

Presented are adjusted odds ratios (aOR) and their 95% confidence intervals (95%CI). All p-values are two-sided. Results are presented separately for maternal and infant healthcare expenditures, adjusted for maternal age, parity, country of origin, parenthood status, and all indicators of SES (i.e. educational level, disposable income, home ownership, and neighborhood poverty)

related to this point is that healthcare expenditures were only available at the annual level. We pragmatically addressed this by only assessing deliveries in December when exploring the association between postpartum care uptake and healthcare expenditure. Although this substantially reduced sample size, statistically significant and clinically relevant associations were still observed.

Our findings stress the need to further explore how equity in care uptake may be promoted. Obstetric healthcare providers should include the social determinants of health in their medical records, and in the referral to postpartum care organizations.<sup>2,15–17</sup> Provision of postpartum care should be tailored according to these determinants to reach poor and other marginalized subpopulations.<sup>18,19</sup> When striving to reduce inequalities in uptake of postpartum care additional determinants, besides those related to a person's SES, should be considered. For example, our results showed that immigrant populations were less likely to receive postpartum care, even when accounting for SES indicators (table 2). This suggests that interventions targeting high-risk groups to increase postpartum care uptake should consider ethnic background in addition to SES-related factors. Cultural factors are likely to explain at least part of this inequity, but this requires further study. Mixed-methods research is needed to assess the facilitators and barriers to receiving postpartum care among low-SES women and those with an immigrant background.

Our results are in line with those observed in other reports; there is a consistent inequity in primary care provision, where more care is provided to the well-off, who need it less, than to the more disadvantaged.<sup>2,4,5,16–18,20</sup> In the Netherlands, a possible barrier to postpartum care uptake is the additional co-payment required for each hour of care. To ensure equitable, universal coverage, policy-makers and health insurers should consider waiving this co-payment, particularly for low SES groups. Furthermore, our study provides evidence to suggest that postpartum care may help reduce subsequent healthcare expenditure, providing an additional incentive for stakeholders to invest in increasing the uptake of care. There is a need to further assess whether explicit resource allocation and priority setting to those in greatest need, perhaps in conjunction with approaches to reduce unnecessary care provision resulting in the over-payment in other sectors may help improve cost-effectiveness of postpartum care provision.

Given the observational nature of this study it is important that findings are reproduced in other settings or using different

methodological approaches (i.e. prospective or randomized studies). Future research should focus on further analysing the (cost-)effectiveness of postpartum care; not only its effectiveness in achieving equity in care provision but also its ability to prevent illness and associated healthcare needs. An in-depth economic evaluation taking into account all expenses made by postpartum care organizations and the potential benefits (e.g. health benefits or value-based health measures) gained by mothers and their children could strengthen a renewed allocation for care provision.

## Supplementary data

Supplementary data are available at *EURPUB* online.

## Acknowledgements

We acknowledge Statistics Netherlands for assistance with project management. We further acknowledge that data are based on the authors' calculations using data files from Statistics Netherlands.

## Funding

This work was supported by ZonMW (The Netherlands Organization for Health Research and Development), research grant number 50-52000-98-173. The study protocol has undergone peer review by the funding body. J.V.B. is supported by personal fellowships from the Netherlands Lung Foundation and Erasmus MC. The funding sources had no role in study design, data collection and analysis, decision to publish or preparation of the article.

*Conflicts of interest:* None declared.

## Key points

- Women of low socioeconomic status are much less likely to receive postpartum care and low uptake of postpartum care is associated with higher health care expenditure after childbirth.
- Substantial inequities in postpartum care provision according to immigration status furthermore exist.
- Interventions to increase postpartum care uptake among those who are most likely to benefit may help reduce health inequalities and subsequent healthcare costs.

## References

- 1 Shaw E, Levitt C, Wong S, et al. Systematic review of the literature on postpartum care: effectiveness of postpartum support to improve maternal parenting, mental health, quality of life, and physical health. *Birth* 2006;33:210–20.
- 2 Marmot M, Friel S, Bell R, et al. Commission on social determinants of H. Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet* 2008;372:1661–9.
- 3 Pillas D, Marmot M, Naicker K, et al. Social inequalities in early childhood health and development: a European-wide systematic review. *Pediatr Res* 2014;76:418–24.
- 4 Ahmed S, Creanga AA, Gillespie DG, Tsui AO. Economic status, education and empowerment: implications for maternal health service utilization in developing countries. *PLoS One* 2010;5:e11190.
- 5 Lindquist A, Kurinczuk JJ, Redshaw M, Knight M. Experiences, utilisation and outcomes of maternity care in England among women from different socio-economic groups: findings from the 2010 National Maternity Survey. *BJOG* 2015;122:1610–7.
- 6 Gray R, Bonellie SR, Chalmers J, et al. Social inequalities in preterm birth in Scotland 1980-2003: findings from an area-based measure of deprivation. *Bjog* 2008;115:82–90.

- 7 Poeran J, Maas AF, Birnie E, et al. Social deprivation and adverse perinatal outcomes among Western and non-Western pregnant women in a Dutch urban population. *Soc Sci Med* 2013;83:42–9.
- 8 Chauvel L, Leist AK. Socioeconomic hierarchy and health gradient in Europe: the role of income inequality and of social origins. *Int J Equity Health* 2015;14:132.
- 9 Currie J, Lin W. Chipping away at health: more on the relationship between income and child health. *Health Aff (Millwood)* 2007;26:331–44.
- 10 Mackenbach JP. The persistence of health inequalities in modern welfare states: the explanation of a paradox. *Soc Sci Med* 2012;75:761–9.
- 11 Benchimol EI, Smeeth L, Guttman A, et al. [The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statement]. *Z Evid Fortbild Qual Gesundheitswes* 2016;115-116:33–48.
- 12 Devillé WW. Herijking stedelijke achterstandsgebieden. Available at: <https://www.nivel.nl/sites/default/files/bestanden/Rapport-Herijking-achterstandsgebieden-2012.pdf>, 2012.
- 13 Autoriteit NZ. Available at: <https://www.nza.nl/english>, 2012.
- 14 Royston P. Multiple imputation of missing values. *Stata J* 2004;3:227–41.
- 15 Gottlieb L, Sandel M, Adler NE. Collecting and applying data on social determinants of health in health care settings. *JAMA Intern Med* 2013;173:1017–20.
- 16 Heiman HA. *Beyond health care: the role of social determinants in promoting health and health equity*. The Henry J Kaiser Family Foundation 2015; Issue brief. Available at: <http://files.kff.org/attachment/issue-brief-beyond-health-care>.
- 17 ACOG Committee Opinion No. 729: Importance of social determinants of health and cultural awareness in the delivery of reproductive health care. *Obstet Gynecol* 2018;131:e43–8.
- 18 Blas E, Gilson L, Kelly MP, et al. Addressing social determinants of health inequities: what can the state and civil society do? *Lancet* 2008;372:1684–9.
- 19 Victora CG, Barreto ML, do Carmo Leal M, et al. Health conditions and health-policy innovations in Brazil: the way forward. *Lancet* 2011;377:2042–53.
- 20 Hart JT. The inverse care law. *Lancet* 1971;1:405–12.

.....  
*The European Journal of Public Health*, Vol. 29, No. 5, 855–861

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Public Health Association. All rights reserved.  
 doi:10.1093/eurpub/ckz089 Advance Access published on 5 June 2019

## Cumulative childhood adversity, adolescent psychiatric disorder and violent offending in young adulthood

Emma Björkenstam<sup>1,2</sup>, Bo Burström<sup>1</sup>, Anders Hjern<sup>3,4</sup>, Bo Vinnerljung<sup>4,5</sup>, Kyriaki Kosidou<sup>6,7</sup>, Lisa Berg<sup>3,8</sup>

1 Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

2 Fielding School of Public Health and California Center for Population Research, Department of Community Health Sciences, University of California, Los Angeles, CA, USA

3 Centre for Health Equity Studies, Stockholm University/Karolinska Institutet, Stockholm, Sweden

4 Clinical Epidemiology/Department of Medicine, Karolinska Institutet, Stockholm, Sweden

5 Department of Social Work, Stockholm University, Stockholm, Sweden

6 Division Public Health Epidemiology, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden

7 Center for Epidemiology and Community Medicine, Stockholm County Council, Stockholm, Sweden

8 Department of Public Health Sciences, Stockholm University, Stockholm, Sweden

**Correspondence:** Emma Björkenstam, Department of Clinical Neuroscience, Karolinska Institutet, SE-171 77 Stockholm, Sweden, Tel: +46 76 212 2704, e-mail: emma.bjorkenstam@ki.se

**Background:** Childhood adversity (CA) is a risk indicator for psychiatric morbidity. Although CA has been linked to violent offending, limited research has considered adolescent psychiatric disorder as a mediating factor. The current study examined whether adolescent psychiatric disorder mediates the association between CA and violent offending. **Methods:** We used a cohort of 476 103 individuals born in 1984–1988 in Sweden. Register-based CAs included parental death, substance abuse and psychiatric disorder, parental criminal offending, parental separation, public assistance, child welfare intervention and residential instability. Adolescent psychiatric disorder was defined as being treated with a psychiatric diagnosis prior to age 20. Estimates of risk of violent offending after age 20 were calculated as incidence rate ratios (IRRs) with 95% confidence intervals (CIs). Mediation was tested with the bootstrap method. **Results:** Exposure to CA was positively associated with violent offending, especially when accumulated. Individuals exposed to 4+ CAs who were also treated for psychiatric disorder had a 12-fold elevated risk for violent offending (adjusted IRR 12.2, 95% CI 10.6–14.0). Corresponding IRR among 4+ CA youth with no psychiatric disorder was 5.1 (95% CI 4.5–5.6). Psychiatric disorder mediated the association between CA and violent offending. **Conclusions:** CA is associated with elevated risk for violent offending in early adulthood, and the association is partly mediated by adolescent psychiatric disorder. Individuals exposed to cumulative CA who also develop adolescent psychopathology should be regarded as a high-risk group for violent offending, by professionals in social and health services that come into contact with this group.

## Introduction

Studies from many western countries have shown that young adults with a history of childhood adversity (CAs) have greater risk for a multitude of problems compared with peers without CAs, especially in the area of psychiatric morbidity.<sup>1–5</sup> CAs generally

include abuse, neglect and growing up in a dysfunctional household, the latter being characterized by substance abuse in the home, mentally ill family members or incarcerated parents.<sup>2</sup>

CAs have also been associated with initiation of criminal behavior.<sup>6–9</sup> Furthermore, research has found that certain